arXiv:math/0409579v2 [math.AP] 25 Feb 2005

EIGENFUNCTIONS FOR PARTIALLY RECTANGULAR BILLIARDS

JEREMY MARZUOLA

1. Introduction

In this note, we further develop the methods of Burq-Zworski [8] to study eigenfunctions for billiards which have rectangular components: these include the Bunimovich billiard, the Sinai billiard, and the recently popular pseudointegrable billiards [2]. The results are an application of a "black box" point of view as presented in [7] by the same authors.



Figure 1. Experimental images of eigenfunctions in a Sinai billiard microwave cavity – see `http://sagar.physics.neu.edu`. We see that there is always a non-vanishing presence near the boundary of the obstacle as predicted by Theorem 3 below.

By a partially rectangular billiard, we mean a connected planar domain, Ω, with a piecewise smooth boundary, which contains a rectangle, $R \subset \Omega$, such that if we decompose the boundary of R, into pairs of parallel segments, $\partial R = \Gamma_1 \cup \Gamma_2$, then $\Gamma_i \subset \partial\Omega$, for at least one i.

We show that for such billiards, the eigenfunctions of the Dirichlet, Neumann, or periodic Laplacian cannot concentrate in closed sets in the interior of the rectangular part. A combination of this elementary result with the now standard, but highly non-elementary, propagation results of Melrose-Sjöstrand [16] and Bardos-Lebeau-Rauch [1], can give further improvements – see [7],[8].

Here, we prove further non-concentration results, away from the obstacle in the Sinai billiard (see Fig.1 and Theorem 3), and along certain trajectories in pseudointegrable billiards, (see Fig.5 and Theorem 4). For recent motivation coming from the study of *quantum chaos* we suggest [2],[8],[10],[17], and references given there.

Acknowledgments. This paper is a development of an unpublished work by N. Burq and M. Zworski, who treated the case of a square billiard rather than a general rectangular billiard. The author is very grateful to Maciej Zworski and Nicolas Burq for allowing me to use results from their unpublished work, as well as many helpful conversations. I would also like to thank Srinivas Sridhar for allowing the use the experimental images shown in Fig.1.

2. SEMICLASSICAL PSEUDODIFFERENTIAL OPERATORS ON A TORUS

In this section, we would like to discuss properties of Pseudodifferential Operators (PDO's) on a torus. To begin, we examine the nature of PDO's and their symbol classes. In $\mathbb{R}^n$, we define a Weyl quantization of an operator $a(x,hD)$ where $a \in \mathcal{S}(\mathbb{R}^{2n}), a = a(x,\xi)$ by:

$$a(x,hD)u(x) = \frac{1}{(2\pi h)^n} \int_{\mathbb{R}^n} \int_{\mathbb{R}^n} e^{\frac{i}{h}\langle x-y,\xi\rangle} a(x,\xi)u(y)dyd\xi,$$

for $u \in \mathcal{S}$ and a symbol class by:

$$S^k_\delta(m) = \{a \in C^\infty(\mathbb{R}^{2n}) \| |\partial^\alpha a| \leq C_\alpha h^{-\delta|\alpha|-k} m \text{ for all multi-indices } \alpha\},$$

where $m : \mathbb{R}^{2n} \to (0,\infty)$ is is an order function, i.e. there exist constants C, N such that

$$m(z) \leq C\langle z-w\rangle^N m(w).$$

We also have

$$S^{-\infty}_\delta(m) = \bigcap_{k=-\infty}^{\infty} S^k_\delta(m).$$

For $k, \delta = 0$ we write simply $S(m)$. On a torus, however, a and all its derivatives are bounded in the x variable, thus for h small and k positive, we need not worry about the derivatives in x, only those in ξ. Also, for k negative, provided that we have the proper local regularity for our symbol a, this definition still works perfectly on a torus.

Note also that we need only work with symbols that are periodic in the x-variable with period determined by the dimensions of the torus. In other words, $a(x,\xi) = a(x+\gamma,\xi)$ for $\gamma \in (a\mathbb{Z})\times(b\mathbb{Z})$, where $a, b \in \mathbb{R}$. Using this relation, we easily see the following propostion.

Proposition 2.1. *If $a(x,\xi)$ is a periodic symbol in x with period γ, then $a(x,D)T_\gamma = T_\gamma a(x,D)$ where $T_\gamma u(x) = u(x-\gamma)$.*

Proof. We easily calculate:

$$a(x,hD)T_\gamma(u) = \int_{\mathbb{R}^n} a(x,\xi)e^{\frac{i}{h}<x-y,\xi>}\hat{u}(y-\gamma)dyd\xi \tag{2.1}$$

$$= \int_{\mathbb{R}^n} a(x-\gamma,\xi)e^{\frac{i}{h}<x-\gamma-\tilde{y},\xi>}\hat{u}(\tilde{y})d\tilde{y}d\xi \tag{2.2}$$

$$= T_\gamma(a(x,D)u) \tag{2.3}$$

□

From the above proposition, it becomes clear that the properties of symbol classes in Euclidean space translate directly to properties of similarly defined symbol classes on a torus. For instance, we have the following result.

Proposition 2.2. *Given* $u(x) = u(x+\gamma)$ *where* γ *is as above, and* u *is* L^2 *on a torus, then* $a(x, hD)u(x)$ *is* L^2 *on the torus, when* $a \in \mathcal{S}_\delta(1)$, $0 \leq \delta \leq \frac{1}{2}$.

Proof. Note that the condition on a implies that it is L^2 bounded. Given a function $u(x)$ which is periodic on a torus, we can write it as $\sum_\gamma T_\gamma u_0$ where $u_0 = \chi(x)u(x)$ and $\chi(x)$ is equal to 1 on a single copy of the torus in the plane and 0 otherwise. Note that no assumptions about the smoothness of $\chi(x)$ are made. Hence, $u_0 \in L^2$ and therefore, so is $a(x, hD)u_0$. Then, $a(x, hD)u(x) = \sum_\gamma T_\gamma a(x, hD)u_0$. The sum converges for each x since $a(x, D)u_0$ will have compact support and we have $a(x, hD)u$ a periodic function that is L^2 on the torus. □

Using similar techniques, we would like to develop the concept of a microlocal defect measure in this setting. As shown in [11], we have the following theorem in Euclidean space:

Theorem 1. *There exists a Radon measure* μ *on* $\mathbb{R}^n$ *and a sequence* $h_j \to 0$ *such that*

$$\langle a^w(x, h_jD)u(h_j), u(h_j)\rangle \to \int_{\mathbb{R}^{2n}} a(x,\xi)d\mu \tag{2.4}$$

for all symbols $a \in S(1)$.

We call μ a microlocal defect measure associated with the family $\{u(h)\}_{0<h\leq h_0}$. Note that an $S(1)$ symbol on the torus corresponds to an $S(1)$ symbol on the plane, therefore this result proves the existence of microlocal defect measures on a torus as well.

Proof. 1. Let $\{a_k\} \in C_c^\infty$ be dense in $C_0(\mathbb{R}^{2n})$. Select a sequence $h_j^1 \to 0$ such that

$$\langle a_1^w(x, h_j^1 D)u(h_j^1), u(h_j^1)\rangle \to \alpha_1.$$

Then, select a subsequence $\{h_j^2\} \subset \{h_j^1\}$ such that

$$\langle a_2^w(x, h_j^2 D)u(h_j^2), u(h_j^2)\rangle \to \alpha_2.$$

Continue such that at the kth step, you take a subsequence $\{h_j^k\} \subset \{h_j^{k-1}\}$ such that

$$\langle a_k^w(x, h_j^k D)u(h_j^k), u(h_j^k)\rangle \to \alpha_k.$$

Then by a diagonal argument, arrive at a sequence h_j such that

$$\langle a_k^w(x, h_j D)u(h_j), u(h_j)\rangle \to \alpha_k$$

for all $k = 1, 2, ...$

2. Define $\Phi(a_k) = \alpha_k$. By a standard theorem on operator norms, we have for each k that

$$|\Phi(a_k)| = |\alpha_k| = \lim_{h_j\to\infty} | < a_k^w u(h_j), u(h_j) > | \leq \limsup_{h_j\to\infty} C\|a_k^w\|_{L^2\to L^2} \leq C \sup |a_k|.$$

The mapping Φ is bounded, linear and densely defined, therefore uniquely extends to a bounded linear functional on $S(1)$, with the estimate

$$|\Phi(a)| \leq C \sup |a|$$

for all $a \in S(1)$. The Riesz Representation Theorem therefore implies the existence of a (possibly complex valued) Radon measure on $\mathbb{R}^{2n}$ such that

$$\Phi(a) = \int_{\mathbb{R}^{2n}} a(x,\xi) d\mu.$$

$\square$

We now quote a general theorem about microlocal defect measures on Euclidean space which we can then apply to a torus. To state the propagation theorem in the form sufficient for our applications, we follow [5].

Let us consider a Riemannian manifold without boundary, M. By partitions of unity we can define semi-classical pseudo-differential operators $a(x, hD_x)$ associated to symbols $a(x,\xi) \in \mathcal{C}^\infty_{\mathrm{c}}(T^*M)$.

Now we consider a sequence (u_n) bounded in $L^2(M)$. satisfying

$$(-h_n^2 \Delta - 1) u_n = 0. \tag{2.5}$$

Using (2.5), as in [12] (see also [5]) we can prove the following result.

Proposition 2.3. *There exist a subsequence (n_k) and a positive Radon measure on T^*M, μ (a semi-classical measure for the sequence (u_n)), such that for any $a \in \mathcal{C}^\infty_{\mathrm{c}}(T^*M)$*

$$\lim_{k \to +\infty} (a^w(x, h_{n_k} D_x) u_{n_k}, u_{n_k})_{L^2(M)} = \langle \mu, a(x,\xi) \rangle. \tag{2.6}$$

Furthermore this measure satisfies

1. *The support of μ is included in the characteristic manifold:*
$$\Sigma \overset{\text{def}}{=} \{(x,\xi) \in T^*M; p(x,\xi) = \|\xi\|_x = 1\} \tag{2.7}$$
where $\|\cdot\|_x$ is the norm for the metric at the point x,
2. *The measure μ is invariant by the bicharacteristic flow (the flow of the Hamilton vector field of p):*
$$H_p \mu = 0, \tag{2.8}$$
3. *For any $\varphi \in \mathcal{C}^\infty_{\mathrm{c}}(T^*M)$,*
$$\lim_{k \to +\infty} \|\varphi u_{n_k}\|^2 = \langle \mu, |\varphi|^2 \rangle. \tag{2.9}$$

The first two properties above are weak forms of the elliptic regularity and propagation of singularities results, whereas the last one states that there is no loss of L^2-mass at infinity in the ξ variable.

Proof. We will prove this proposition only for the case of a torus, but the methods are applicable to any manifold.

(0) (Positivity) We need to show that $a \geq 0$ implies

$$\int_{\mathbb{T}^2 \times \mathbb{R}^2} a(x,\xi) d\mu \geq 0.$$

Since $a \geq 0$, using the Garding inequality, we see that:

$$a^w(x, hD) \geq -Ch.$$

Let $h = h_j \to 0$, to see:

$$\int_{\mathbb{T}^2 \times \mathbb{R}^2} a d\mu = \lim_{j \to \infty} \langle a^w(x, h_j D) u(h_j), u(h_j) \rangle \geq 0.$$

(1) (Support of μ) Let a be a smooth function such that $\mathrm{supp}(a) \cap p^{-1}(1) = \emptyset$. We must show

$$\int_{\mathbb{T}^2 \times \mathbb{R}^2} a d\mu = 0.$$

Select $\chi \in C_c^\infty(\mathbb{T}^2 \times \mathbb{R}^2)$ such that

$$\mathrm{supp}(a) \cap \mathrm{supp}(\chi) = 0.$$

Then,

$$a^w(x, hD) \left(((p-1)^w + i\chi^w)^{-1} (p-1)^w \right)(x, hD) = a^w(x, hD) + \mathcal{O}(h^\infty)_{L^2 \to L^2}.$$

Apply $a^w(x, hD)$ to $u(h)$ to see that $a^w(x, hD)u(h) = o(1)$ and thus $\langle a^w(x, hD)u(h), u(h)\rangle \to 0$. But,

$$\langle a^w(x, hD)u(h_j), u(h_j)\rangle \to \int_{\mathbb{T}^2 \times \mathbb{R}^2} a d\mu.$$

(2) (Flow Invariance) Select a as above, then

$$\begin{aligned} \langle [p^w, a^w]u(h), u(h)\rangle &= \langle (p^w a^w - a^w p^w)u(h), u(h)\rangle && (2.10)\\ &= \langle a^w u(h), p^w u(h)\rangle - \langle p^w u(h), (a^w)^* u(h)\rangle && (2.11)\\ &= o(h) \text{ as } h \to 0. && (2.12) \end{aligned}$$

However, $[p^w, a^w] = \frac{h}{i}\{p^w, a^w\} + O(h^2)$. Hence,

$$\langle [p^w, a^w]u(h), u(h)\rangle = \frac{h}{i}\langle \{p, a\}^w u(h), u(h)\rangle + \langle o(h)u(h), u(h)\rangle.$$

As we let $h_j \to 0$, we get:

$$\int_{\mathbb{T}^2 \times \mathbb{R}^2} \{p, a\} d\mu = 0.$$

So, if Φ_t is the flow generated by the Hamiltonian vector field H_p, then

$$\frac{d}{dt}\int_{\mathbb{T}^2 \times \mathbb{R}^2} (\Phi_t * a) d\mu = \int_{\mathbb{T}^2 \times \mathbb{R}^2} (H_p a)(\Phi_t) d\mu = \int_{\mathbb{T}^2 \times \mathbb{R}^2} \{p, a\} d\mu = 0.$$

Now, (3) follows easily by looking at the operator $|\varphi(x, \xi)|^2$ and applying the result about existence of a microlocal defect measure. ☐

3. Partially rectangular billiards

In this section we will need to recall the basic control results [4],[7] for rectagles, and the propagation results [16],[1],[5],[6] for billiards. Since in the specific application presented in Section 4 we only use propagation away from the boundary, that is the only case we will review.

The following result from [4] is related to some earlier control results of Haraux [13] and Jaffard [14].[1]

Proposition 3.1. *Let Δ be the Dirichlet, Neumann, or periodic Laplace operator on the rectangle $R = [0, 1]_x \times [0, a]_y$. Let ω_x be a non-empty open subset of $[0, 1]$. Then for any non-empty $\omega \subset R$ of the form $\omega = \omega_x \times [0, a]_y$, there exists C such that for any solutions of*

$$(\Delta - z)u = f \quad \text{on } R, \ u\restriction_{\partial R} = 0 \tag{3.1}$$

we have

$$\|u\|^2_{L^2(R)} \le C\left(\|f\|^2_{H^{-1}([0,1]_x; L^2([0,a]_y))} + \|u\restriction_\omega\|^2_{L^2(\omega)} \right) \tag{3.2}$$

---

[1] We remark that as noted in [4] the result holds for any product manifold $M = M_x \times M_y$, and the proof is essentially the same.

Proof. We will consider the Dirichlet case (the proof is the same in the other two cases) and decompose u, f in terms of the basis of $L^2([0,a])$ formed by the Dirichlet eigenfunctions $e_k(y) = \sqrt{2/a}\sin(2k\pi y/a)$,

$$u(x,y) = \sum_k e_k(y) u_k(x), \qquad f(x,y) = \sum_k e_k(y) f_k(x). \tag{3.3}$$

We get for u_k, f_k the equation

$$\left(\Delta_x - \left(z + (2k\pi/a)^2\right)\right) u_k = f_k, \qquad u_k(0) = u_k(1) = 0. \tag{3.4}$$

We now claim that

$$\|u_k\|^2_{L^2([0,1]_x)} \leq C\left(\|f_k\|^2_{H^{-1}([0,1]_x)} + \|u_k\restriction_{\omega_x}\|^2_{L^2(\omega)}\right), \tag{3.5}$$

from which, by summing the squares in k, we get (3.2).

To see (3.5) we can use the propagation result above 2.3 in dimension one, but in this case an elementary calculation is easily available – see [8]. ☐

The following theorem is an easy consequence of Proposition 3.1:

Theorem 2. *Let Ω be a partially rectangular billiard with the rectangular part $R \subset \Omega$, $\partial R = \Gamma_1 \cup \Gamma_2$, a decomposition into parallel components satisfying $\Gamma_2 \subset \partial\Omega$. Let Δ be the Dirichlet or Neumann Laplacian on Ω. Then for* any *neighbourhood of Γ_1 in Ω, V, there exists C such that*

$$-\Delta u = \lambda u \implies \int_V |u(x)|^2 dx \geq \frac{1}{C}\int_R |u(x)|^2 dx\,, \tag{3.6}$$

that is, no eigenfuction can concentrate in R and away from Γ_1.

Proof. Let us take x, y as the coordinates on the stadium, so that x parametrizes $\Gamma_2 \subset \partial\Omega$ and y parametrizes Γ_1, then

$$R = [0,1]_x \times [0,a]_y\,.$$

Let $\chi \in \mathcal{C}^\infty_{\rm c}((0,1))$ be equal to 1 on $[\varepsilon, 1-\varepsilon]$. Then $\chi(x)u(x,y)$ is a solution of

$$(\Delta - z)\chi u = [\Delta, \chi]u \text{ in } R \tag{3.7}$$

with the boundary conditions satisfied on ∂R. Applying Proposition 3.1, we get

$$\|\chi u\|_{L^2(R)} \leq C\left\|[\Delta,\chi]u\|_{H_x^{-1};L_y^2} + \|u\restriction_{\omega_\varepsilon}\|_{L^2(\omega_\varepsilon)}\right) \leq C'\|u\restriction_{\omega_\varepsilon}\|_{L^2(\omega_\varepsilon)}\,, \tag{3.8}$$

where ω_ε is a neighbourhood of the support of $\nabla\chi$. Since a neighbourhood of Γ_1 in Ω has to contain ω_ε for some ε, (3.6) follows. ☐

4. APPLICATIONS

In [7] and [8], Proposition 3.1 is used to prove that in the case of the Bunimovich billiard shown in Fig.2, the states have nonvanishing density near the vertical boundaries of the rectangle. That follows from Theorem 2 which shows that we must have positive density in the wings of the billiard, and the propagation result (in the boundary case) based on the fact that any diagonal controls a disc geometrically (see [7, Section 6.1]; in fact we can use other control regions as shown in Fig.2). Here we consider another case which accidentally generalizes a control theory result of Jaffard [14].

The Sinai billiard (see Fig.1) is defined by removing a strictly convex open set, $\mathcal{O}$, with a $\mathcal{C}^\infty$ boundary, from a flat torus, $\mathbb{T}^2_{a,b} \overset{\rm def}{=} (a\mathbb{S}^1) \times (b\mathbb{S}^1)$:

$$S \overset{\rm def}{=} \mathbb{T}^2_{a,b} \setminus \mathcal{O}\,.$$



Figure 2. Control regions in which eigenfunctions have positive mass and the rectangular part for the Bunimovich stadium.

The following theorem results by applying Theorem 1 to a torus with sides of arbitrary length.

Theorem 3. *Let V be any open neighbourhood of the convex boundary, $\partial\mathcal{O}$, in a Sinai billiard, S. If Δ is the Dirichlet or Neumann Laplace operator on S then there exists a constant, $C = C(V)$, such that*

$$-h^2\Delta u = E(h)u \implies \int_V |u(x)|^2 dx \geq \frac{1}{C}\int_S |u(x)|^2 dx\,, \tag{4.1}$$

for any h and $|E(h) - 1| < \frac{1}{2}$.

Proof. First note that we can easily limit ourselves to the case where our flat torus has one side of length 1 and one side of length a. Suppose that the result is not true, in other words, there exists a sequence of eigenfunctions u_n, $\|u_n\| = 1$, with the corresponding eigenvalues $\lambda_n \to \infty$, such that $\int_V |u_n(x)|^2 dx \to 0$.

We first observe that the only directions in the support of the corresponding semi-classical defect measure, μ, have to be "rational", in other words, the trajectory must travel along a line of slope $\frac{ma}{n}$ where $m, n \in \mathbb{N}$. The projection of a trajectory with an irrational direction is dense on the torus and hence must encounter the obstacle $\partial\mathcal{O}$ (and consequently V). The propagation result recalled in Proposition 2.3, part (3), gives a contradiction by choosing a proper test function ϕ which is nonzero on the support of the measure μ resulting from our sequence of eigenfunctions (remark that we apply this result as long as the trajectory does not encounter the obstacle and consequently we need only the *interior* propagation).

Hence let us assume that there exists a rational direction in the support of the measure which then contains the periodic trajectory in that direction. As shown in Fig. 3 we can find a maximal rectangular neighbourhood of the projection of that trajectory which avoids the obstacle.

The rectangle can be described as $R = [0, a_1]_{x_1} \times [0, b_1]_{y_1}$ with the the y_1 coordinate parametrizing the trajectory. Let ϵ, $\delta > 0$ be small. Let u be an eigenfunction in our sequence and define

$$\chi = \{\chi(x_1) \in \mathcal{C}^\infty_{\rm c}(\mathbb{T}^2) \mid \chi(x_1) = 1 \text{ for all } x_1 \in (\epsilon, a_1 - \epsilon),\ \chi(x_1) = 0 \text{ outside } (0, a_1)\}.$$

Note that we can then write $\chi = \chi(x, y)$ for $(x, y) \in \mathbb{T}^2_{1,a}$ as x_1 is simply a rotation and translation of the standard coordinates. Then $\chi(x, y)u(x, y)$ is a function on all of R satisfying the periodicity condition. Let $\Phi_\xi(\nu) = \Phi(\xi - \nu)$, where we define

$$\Phi = \{\Phi(\xi) \in \mathcal{C}^\infty_{\rm c}(\mathbb{R}^2) \mid \Phi(\xi) = 1 \text{ for } \xi \in B(0, \delta),\ \Phi(\xi) = 0 \text{ for } \xi \in \mathbb{R}^2 \setminus B(0, 2\delta)\},$$

where $B(0, \delta)$ is a ball centered at 0 of radius δ. Note, due to the compact support of this function in ξ, $\Phi_\xi(D)$ is in the symbol class $S(\langle\xi\rangle^{-N})$ for any N. Let Δ_R is the (periodic) Laplacian on R. Using Fourier decomposition we can arrange that $[\Delta_R, \Phi_\xi(D)] = 0$. Since our eigenfunction



FIGURE 3. A maximal rectangle in a rational direction, avoiding the obstacle. Because the parrallelogram is certainly periodic and our region has uniform width, it is clear that the resulting rectangle is periodic.

is only defined on $\mathbb{T}^2_{1,a} \setminus \mathcal{O}$, let us introduce a smooth function χ_0 which is 0 on a neighborhood U of the obstacle and 1 on $\mathbb{T}^2_{1,a} \setminus V$ where $U \subset V$. Choose U and V such that $\chi\chi_0 = \chi$. Hence,

$$(-h^2\Delta_R - E(h))\Phi_\xi(D)\chi\chi_0 u = [-h^2\Delta_R, \Phi_\xi(D)\chi]u = \Phi_\xi(D)[-h^2\Delta_R, \chi]\chi_0 u + \mathcal{O}(h^\infty)\,, \quad \|u\| = 1\,,$$

by the properties of $S(\langle\xi\rangle^{-N})$ operators acting on L^2 functions on a torus. As in the proof of Proposition 3.1, we now see that

$$\|\Phi_\xi \chi\chi_0 u\|_{L^2} \leq C \int_\omega |\chi_0 u|^2 + \mathcal{O}(h^\infty)\,, \tag{4.2}$$

where ω is a neighbourhood of $\nabla\chi$ (in the calculus of semi-classical pseudo-differential operators). Since the semi-classical defect measure of $\Phi_\xi\chi\chi_0 u$ (which is $|\Phi_\xi\chi\chi_0|^2 \times \mu$) was assumed to be non-zero, (4.2) shows that the measure of $\chi_0 u\restriction_\omega$ is non zero and consequently there is a point in the intersection of the supports of μ and $\chi_0 u\restriction_\omega$. But μ is invariant by the flow (as long as it does not intersect the obstacle) and hence, once we choose ϵ, δ small enough such that all the cut-offs above are very close to the boundary of R, its support can be made intersect any neighbourhood of $\partial\mathcal{O}$. $\square$

Now, from the above theorem, we see the following simple, but important consequences:

Remark 1. Let $S = \mathbb{T}^2_{a,b} \setminus \mathcal{O}$ where $\mathcal{O}$ is sufficiently smooth in the case of Neumann boundary conditions, but otherwise lacking restrictions. Then, for V any open neighborhood of $\partial\mathcal{O}$, and u a solution of $-h^2\Delta u = E(h)u$ as above, then (4.1) is satisfied. This follows from the above argument as the convexity of the obstacle was never used. Thus, the result holds for any obstacle (even connectedness is not assumed here) and is applicable to the special case of pseudointegrable billiards (see for instance [2] for motivation and description). In the next section, we use an argument similar to that above in order to say even more about concentration along trajectories in specific pseudointegrable billiards. By an elementary reflection principle, the result also holds for an obstacle inside a square with Dirichlet or Neumann conditions on the boundary of the square.

Remark 2. The proof above gives in fact the following estimate for any open neighbourhood, say V, of the obstacle:

$$\begin{gathered} \exists C; \forall u,\ f \in L^2(S) \text{ solutions of } (-\Delta + \lambda)u = f, \qquad u\restriction_{\partial S} = 0 \\ \|u\|_{L^2(S)} \leq C\left(\|f\|_{L^2(S)} + \|u\mathbb{1}_V\|_{L^2(V)}\right) \end{gathered} \tag{4.3}$$

and according to [7, Theorem 4], this implies that the Schrödinger equation in S is exactly controllable by V in finite time. In fact, by working on the time evolution equation, we could strengthen this result allowing an arbitrarily small time.

This latter result was previously known [14] for the particular case $\Theta = \emptyset$ ($S = \mathbb{T}^2$) but the proof was based on subtle results about Fourier series [15].

Remark 3. As shown in [7, Theorem 2′], the results of Ikawa and Gérard on scattering by two convex obstacles (see [7] and references given there) give an estimate on the maximal concentration of an eigenfunction (or a quasimode) on a closed orbit in a Sinai billiard. Let $\chi \in \mathcal{C}^\infty(S; [0,1])$ be supported in a small neighbourhood of a closed transversally reflecting orbit. Then for any family $(-\Delta - \lambda)u_\lambda = \mathcal{O}(\lambda^{-\infty})$, $\|u_\lambda\| = 1$,

$$C \int_S |u(x)|^2 (1 - \chi(x)) dx \geq \frac{1}{\log \lambda},$$

that is a concentration on a closed trajectory, if at all possible, has to be very weak.

5. Pseudointegrable Billiards

We define a pseudointegrable billiard to be a plane polygonal billiard with corners whose angles are of the form $\frac{\pi}{n}$, for any integer n (see [3]). In particular, we will be working with the billiard $P = \mathbb{T}^2_{a,b} \setminus S$ where S is a slit that is parrallel to a side of the torus but not a closed loop. In Remark 1, we point out that Theorem 3 allows us to make statements about the L^2 mass of eigenfunctions in a neighborhood of the slit for pseudointegrable billiards. For this particular type of billiard, it would be ideal to state that every eigenfunction must have non-zero mass in a small neighborhood of the edges of the slit (see Fig. 4). In this section, we prove a weaker result about non-concentration along certain classical trajectories in P of semiclassical defect measures obtained from eigenfunctions u such that $(-\lambda - \Delta)u = 0$ on P.

As with the Sinai billiard, the classical behavior of trajectories must be taken into account in our treatment of this problem. There cannot be concentration along trajectories that do not hit the slit as shown by Theorem 3. If a trajectory has irrational slope, it is dense in P, and thus has mass near the edges of the slit as in Section 4. Therefore, for our purpose, we concern ourselves only with rational trajectories which intersect the slit at some point. As we are dealing with periodic boundary conditions, let us consider the plane tiled with copies of the billiard P.



FIGURE 4. A pseudointegrable billiard P consisting of a torus with a slit, S along which we have Dirichlet boundary conditions. We would like to show that eigenfunctions of the Laplacian on this torus must have concentration in the shaded regions V_1 and V_2.

Assume that P is oriented such that S is parralel to the y-axis. Let $\gamma \in S^*(P)$ be a trajectory. Given the natural projection

$$\pi_1 : S^*(P) \to P,$$

we take $\gamma' = \pi_1(\gamma)$, or the physical path along which the trajectory travels. Consider the projection

$$\tilde{\pi} : \mathbb{R}^2 \to \mathbb{T}^2_{a,b}.$$

We see that

$$\tilde{\pi} : \mathbb{R}^2 \setminus \tilde{S} \to P, \text{ where } \tilde{S} = \tilde{\pi}^{-1}(S).$$

Define

$$\pi_2 : S^*(\mathbb{R}^2 \setminus \tilde{S}) \to S^*(P)$$

to be the obvious projection. Let $\tilde{\gamma} = \pi_2^{-1}(\gamma)$. We can write

$$\tilde{\gamma} = \cup_{j=1}^{\infty} \gamma_j,$$

where each γ_j is a trajectory in $S^*(\mathbb{R}^2 \setminus \tilde{S})$. We note that by construction, $\gamma_i \cap \gamma_j = \emptyset$ for $i \neq j$. To see this, assume that $\gamma_i \cap \gamma_j = (x, \xi)$. Then, $\gamma_i = \gamma_j$ as they would represent trajectories which travel through the same point in the same direction. Now, let

$$\pi_1^* : S^*(\mathbb{R}^2 \setminus \tilde{S}) \to \mathbb{R}^2 \setminus \tilde{S}.$$

Select one trajectory from the above union, say γ_1. Let $\gamma_1' = \pi_1^*(\gamma_1)$. We see that either γ_1' is bounded in the x-direction or γ_1' is unbounded in the x-direction. Note that this property then holds for all γ_j, $j \in \mathbb{N}$. For a trajectory γ, if the resulting path γ_1' is bounded in the x-direction, we say γ is x-bounded. We define γ as x-unbounded if γ_1' is unbounded in the x-direction. See Fig. 5 for examples. Now, we are prepared to state our theorem concerning the billiard P.

Theorem 4. *Let γ be an x-bounded trajectory on $P = \mathbb{T}^2 \setminus S$. If Δ is the Dirichlet Laplace operator on P then there exists no microlocal defect measure obtained from the eigenfunctions on P such that* $\text{supp}\ (d\mu) = \gamma$.



a. b.

c. d.

Above, a. and b. represent typical x-bounded trajectories, while c. and d. represent x-unbounded trajectories.

FIGURE 5. Some examples of x-bounded and x-unbounded trajectories.

Proof. Let γ' be as above. Let V_ϵ be an ϵ neighborhood of γ'. If the theorem were false, we would have a sequence of eigenfunctions u_n, $\|u_n\|_{L^2} = 1$ with the property

$$\int_{P \setminus V_\epsilon} |u_n|^2 dx \to 0,$$

for any ϵ. We show that this is impossible.

For each u_n, we have $(-\Delta - \lambda_n)u_n = 0$, $u_n|_S = 0$, $u_n \in L^2(P)$. Let $\tilde{\pi}$ be as above. We define the sequence $\tilde{u}_n = \tilde{\pi}^{-1}u_n$. We have $(-\Delta - \lambda_n)\tilde{u}_n = 0$, $\tilde{u}_n|_{\tilde{S}} = 0$, and $\tilde{u}_n \in L^2_{\text{per}}(\mathbb{R}^2 \setminus \tilde{S})$.

If $\pi_2 : S^*(\mathbb{R}^2 \setminus \tilde{S}) \to S^*(P)$ is as above and $\tilde{\gamma} = \pi_2^{-1}(\gamma)$, then $\tilde{u}_n \to d\tilde{\mu}$ with

$$\text{supp } (d\tilde{\mu}) = \tilde{\gamma} \subset S^*(\mathbb{R}^2 \setminus \tilde{S})\,.$$

Now, let $\pi_1^* : S^*(\mathbb{R}^2 \setminus \tilde{S}) \to \mathbb{R}^2 \setminus \tilde{S}$ be as above. Select one trajectory, say γ_1. As γ_1 is x-bounded, $\gamma_1' = \pi_1^*(\gamma_1)$ is contained in a strip in the plane which is infinite in the y-direction and bounded in the x-direction. Thus, γ_1' is contained in a strip, C_0, with minimal width in the

x-direction. Then, $\tilde{u}_n$ satisfies $(-\Delta-\lambda_n)\tilde{u}_n = 0$ on C_0, is periodic in the y-direction, and satisfies the following boundary conditions in the x-direction: Dirichlet boundary conditions along the slits that intersect the boundary of C_0 and periodic boundary conditions otherwise.

Without loss of generality, we can choose the x-coordinates such that the boundaries of C_0 are $x = -R$ and $x = 0$. We can then reflect to a strip, say $\tilde{C}_1$, with boundaries $x = -R$ and $x = R$, by defining a new function on $\tilde{C}_1$ by:

$$\tilde{u}_n^{(1)}(x,y) = \begin{cases} \tilde{u}_n(x,y) & x \in [-R,0], \\ -\tilde{u}_n(-x,y) & x \in (0,R). \end{cases}$$

Note that $\tilde{u}_n$ is periodic with period $2R$. As a result, we have

$$(-\Delta - \lambda_n)\tilde{u}_n^{(1)} = f_n^{(1)}$$

on $\tilde{C}_1$, where

$$f_n^{(1)} = 2u(0,y)\delta_0'(x) - 2u(R,y)\delta_R'(x).$$

We note that $f_n^{(1)}$ is supported away from the slits, $\tilde{S}$.

Define

$$\pi_1^\sharp(x,y) = \begin{cases} (x,y) & -R \leq x \leq 0, \\ (-x,y) & 0 \leq x \leq R. \end{cases}$$

If $\pi_1^\sharp \ : \ \tilde{C}_1 \to C_0$, then

$$(\pi_1^\sharp)^{-1}(\bigcup_j \gamma_j')$$

is again a union of paths resulting from disjoint trajectories. Now, we iterate this procedure a finite number of times, stopping the iteration when the disjoint trajectories in the lift intersect each slit only once.

After each reflection, we restrict to a new minimal width strip, say C_i. Let us call $\tilde{C}_i$ the strip resulting from the ith reflection. We define $\pi_i^\sharp : \tilde{C}_i \to C_{i-1}$ for $1 \leq i < N$ such that

$$\pi_i^\sharp(x,y) = \begin{cases} (x,y) & (x,y) \in C_{i-1}, \\ (2R_i - x, y) & (x,y) \in C_{i-1}'. \end{cases}$$

Here, C_{i-1}' is defined as the reflected strip and $x = R_{i-1}$ is the line of reflection for $\tilde{C}_i$. We can subsequently define $f_n^{(i)}$ as a sum of delta functions resulting from jumps that occur after reflection, similar to $f_n^{(1)}$ above. We also have $\pi^N : \mathbb{R}^2 \to C_N$, the obvious projection that results after we tile the plane with copies of C_N. So, we have:

$$\mathbb{R}^2 \overset{\pi_N}{\to} C_N \subset \tilde{C}_N \overset{\pi_N^\sharp}{\to} C_{N-1} \subset \tilde{C}_{N-1} \overset{\pi_{N-1}^\sharp}{\to} \ldots \overset{\pi_2^\sharp}{\to} C_1 \subset \tilde{C}_1 \overset{\pi_1^\sharp}{\to} C.$$

Note that

$$\pi_N^{-1}(\gamma_1') = \bigcup_j \gamma_{1,j}',$$

where $\{\gamma_{1,j}'\}$ is the set of all paths in C_N generated by the trajectory γ_1 and the periodicity in y.

After a finite number of reflections, we "unfolded" γ_1' to be a periodic line on a large strip, C_N, which does not intersect a slit anywhere. Now, let us choose Φ_ξ, χ, and χ_0 as above in order to cut-off microlocally on this strip around γ_1. Again, recall that we can set $\chi\chi_0 = \chi$. As $f_n^{(i)}$ is supported only in between the slits for each $i \in \mathbb{N}$, $1 \leq i \leq N$, by choosing Φ_ξ to commute with the periodic Laplacian, we have

$$(-h^2\Delta_R - E(h))\Phi_\xi\chi\chi_0 u_n = \Phi_\xi\chi f_n + [-h^2\Delta_R, \Phi_\xi\chi]u = \Phi_\xi[-h^2\Delta_R, \chi]\chi_0 u + \mathcal{O}(h^\infty)\,, \quad \|u\| = 1\,.$$

Thus, the result follows by contradiction from the proof of Theorem 3. $\square$



In bold, we haveγ_1 in the plane.
The union of all trajectories here gives $\tilde{\gamma}$.

In bold, we have γ_1 in C_0

γ_1 in the strip $\tilde{C}_1$. The inhomogeneity resulting from reflection will be supported along the blue lines. Note also that we have elected to show only γ_1 for simplicity.



γ_1 in C_2 after another reflection.

γ_1 as a periodic trajectory in $\mathbb{R}^2$ after a final reflection and multiplication by a microlocal cut-off function.

Figure 6. This diagram describes how we "unfold" the eigenfunctions in order to derive a contradiction.

Remark 4. Though this result only shows non-concentration, the proof of Theorem 3 can be used to show if γ is an x-bounded trajectory and u is an eigenfunction supported on $\gamma' = \pi_1(\gamma)$, then in fact there must be mass at the edges of the slits as desired.

Remark 5. If instead of a torus, we had Dirichlet boundary conditions on the boundary of the rectangle as well as the slit, then this non-concentration result can also be applied by an elementary reflection principle argument.

References

[1] C. Bardos, G. Lebeau and J. Rauch. Sharp sufficient conditions for the observation, control, and stabilization of waves from the boundary. *SIAM J. Control Optim.* 30:1024–1065, 1992.

[2] E. Bogomolny, U. Gerland, and C. Schmit. Models of intermediate spectral statistics, *Phys. Rev. E* 59:1315-1318, 1999.

[3] E. Bogomolny and C. Schmit. Structure of Wave Functions of Pseudointegrable Billiards, *Phys. Rev. Lett.* 92:244102, 2004.

[4] N. Burq. Control for Schrodinger equations on product manifolds. *Unpublished*, 1992

[5] N. Burq. Semi-classical estimates for the resolvent in non trapping geometries. *Int. Math. Res. Notices*, 5:221–241, 2002.

[6] N. Burq and P. Gérard, Condition nécessaire et suffisante pour la contrôlabilité exacte des ondes. *Comptes Rendus de L'Académie des Sciences*, 749–752,t.325, Série I, 1996

[7] N. Burq and M. Zworski. Geometric control in the presence of a black box. *JAMS*, to appear.

[8] N. Burq and M. Zworski. Bouncing ball modes and quantum chaos. *SIAM Review*, to appear.

[9] N. Burq and G.Lebeau. Mesures de défaut de compacité, application au système de Lamé. *Ann. Sci. École Norm. Sup. (4)*, No 34, 817-870, 2001.

[10] H. Donnelly. Quantum unique ergodicity. *Proc. Amer. Math. Soc.* 131:2945-2951, 2003.

[11] L. C. Evans and M. Zworski. Lectures on Semiclassical Analysis. *Unpublished Lecture Series*, 2003.

[12] P. Gérard and E. Leichtnam. Ergodic Properties of Eigenfunctions for the Dirichlet Problem. *Duke Mathematical Journal,* No 71, 559–607, 1993

[13] A. Haraux. Séries lacunaires et contrôle semi-interne des vibrations d'une plaque rectangulaire. *J. Math. Pures Appl.* 68-4:457–465, 1989.

[14] S. Jaffard. Contrôle interne exact des vibrations d'une plaque rectangulaire. *Portugal. Math.* 47 (1990), no. 4, 423-429.

[15] J.P. Kahane. Pseudo-périodicité et séries de Fourier lacunaires. *Annales Sc. de l'Ecole Normale Supérieure* 79, 1962.

[16] R.B. Melrose and J. Sjöstrand. Singularities of Boundary Value Problems I & II, *Communications in Pure Applied Mathematics*, 31 & 35, 593- 617 & 129-168, 1978 & 1982.

[17] S. Zelditch. Quantum unique ergodicity. `math-ph/0301035`

Mathematics Department, University of California, Evans Hall, Berkeley, CA 94720, USA
E-mail address: `marzuola@math.berkeley.edu`